

Clinical Study Design


- **Open label, observational study**
- **Inclusion**
 - Male and female (Adult)
 - History of allergic rhinitis (≥ 1 year)
 - Positive skin prick test (RAST) to *Parietaria officinalis* pollen
 - Baseline nasal and/or ocular symptoms of seasonal rhinitis
- **Dosing:** 2 tablets per day. In the morning and evening at or after mealtime
- **Treatment period:** 30 day duration
- **Data collection:** Baseline and D30

Evaluation Methods

- **Evaluation methods:**
 - Relief of signs and symptoms of ARC using the Total Symptoms Score and Scale (TSS):
 - 6 clinical parameters graded from 0-3 based on episodes per day
 - 0 = absence
 - 1 = 1-5 episodes
 - 2 = 6-10 episodes
 - 3 = ≥ 11 episodes
 - Reduction of the consumption of anti-allergic drugs (patient recorded at baseline and D30)
- **Safety:** Monitoring adverse events (AEs)

AllerVarx™ Efficacy


Reduction of the average of the scores of individual symptoms measured at baseline and after 30D of AllerVarx™ use



Highly significant reduction ($p < 0.0001$) of each individual symptom compared to baseline

AllerVarx™ Efficacy

Reduction of the average score of the Total Symptom Score Scale at baseline and after 30 days of AllerVarx™ use



Gender	Symptom score reduction
Females	72%
Males	68%

Highly significant reduction in Total Symptom Scores of the overall symptoms by approximately 70%

AllerVarx™ Efficacy

Reduction of consumption of anti-allergic drugs by gender

	Female (n=16)	Male (n=7)
Average Age	44	46
Reduction of anti-allergic drugs	76%	67%

Reduction of use of antihistamines by 73% by end of study

Efficacy & Safety Summary

- **AllerVarx™ containing quercetin, perilla dry extract and vitamin D₃, is effective in reducing nasal and / or ocular symptoms in subjects with seasonal allergic rhinitis**
 - TSS: highly significant reduction ($p > 0.001$) of the overall symptoms, with a reduction of approximately 70%
 - Individual Symptoms: highly significant reduction ($p > 0.001$) of each symptom
 - Reduction of use of anti-allergic drugs (73%)
- **Well tolerated**
 - No side effects recorded
 - All subjects completed the study